arXiv:astro-ph/9710281v1 24 Oct 1997

# LARGE GROUPS IN THE CHILE-UK QUASAR SURVEY

PETER R. NEWMAN, ROGER G. CLOWES
*Centre for Astrophysics, University of Central Lancashire,*
*Preston PR1 2HE, U.K.*

LUIS E. CAMPUSANO
*Observatorio Astronómico Cerro Calán, Universidad de Chile,*
*Casilla 36D, Santiago, Chile*

MATTHEW J. GRAHAM
*Centre for Astrophysics, University of Central Lancashire,*
*Preston PR1 2HE, U.K.*

The Chile-UK quasar survey, a new-generation 140 deg$^2$ UVX survey to $B = 20$, is now $\sim 25\%$ complete. The catalogue currently contains 319 quasars and 93 emission line galaxies. Using the minimal-spanning tree method, we have independently confirmed the $\sim 200\,h^{-1}$ Mpc group of quasars at $z \simeq 1.3$ discovered by Clowes & Campusano (1991). We have discovered a new $\sim 150\,h^{-1}$ Mpc group of 13 quasars at median $z \simeq 1.51$. The null hypothesis of a uniform, random distribution is rejected at a level of significance of 0.003 for both groups.

## 1 The Chile-UK quasar survey

We are conducting an ultraviolet-excess (UVX) quasar survey using the 2°-field, 128-fibre, multi-object spectrograph on the du Pont 2.5-m telescope at Las Campanas, to study large quasar groups and the general clustering properties of quasars for $0.4 \lesssim z \lesssim 2.2$. The survey includes the largest structure previously known: the $\sim 200\,h^{-1}$ Mpc group of $\geq 18$ quasars at $z \simeq 1.3$ (the CC91 group [1,2]). Survey candidates are all objects with $(U - B) \leq -0.3$ and $16 \leq B \leq 20$ on UK Schmidt Telescope plates digitized by SuperCOSMOS.

## 2 Minimal spanning tree identification of large groups

We used the MST method [3] to identify large groups in the quasar catalogue, and to determine their statistical significance. The critical separation distance which maximizes the number of groups of connected quasars is $44\,h^{-1}$ Mpc. Figure 1 shows the two most significant groups. Group A, 6 quasars at median $z \simeq 1.24$ coincides with the core of the CC91 group,[4] independently confirming this group. Group B, 13 quasars at median $z \simeq 1.51$ spanning $\sim 150\,h^{-1}$ Mpc, is a new detection. These results suggest that parallel sheets or cellular structure and voids existed by $z \simeq 1.5$, and may be related to recent reports of



Figure 1: The main survey area (shaded) and the area nominally observed so far (unshaded – the actual area observed is slightly larger). Dots (labelled with redshifts) show the survey quasars in MST groups A (left) and B (right). Lines show 2D projections of the 3D MST edges, each shorter than the critical distance. Crosses mark other UVX quasars in the same redshift range but at larger separations. In the left panel, open circles show the core MST group of 6 quasars from the CC91 objective-prism survey. Three quasars are common to the CC91 and UVX groups. The other three CC91 quasars not in the UVX group are within the critical distance of the UVX group, but are too faint to be in the UVX sample.

large-scale structure from observations of the galaxy distribution and of gas at high redshift.

## Acknowledgments

We thank the Carnegie Institution of Washington and CTIO for telescope time. PRN is supported by a UK PPARC research studentship. LEC was partially supported by FONDECYT grant 1970735.

## References

1. R.G. Clowes, L.E. Campusano, *MNRAS* **249**, 218 (1991).
2. R.G. Clowes, L.E. Campusano, M.J. Graham in *Wide Field Spectroscopy and the Distant Universe*, eds. S.J. Maddox and A. Aragón–Salamanca (World Scientific, Singapore, 1995), p. 400.
3. M.J. Graham, R.G. Clowes, L.E. Campusano, *MNRAS* **275**, 790 (1995).
4. M.J. Graham, PhD thesis, University of Central Lancashire (1997).